



Howard Johnson's

# Cuatro with Dimension® plus Quinclorac

### ACTIVE INGREDIENT:

2,4-D	.....0.675%
Dicamba	.....0.060%
Dithiopyr	.....0.121%
Quinclorac	.....0.346%

OTHER INGREDIENTS:	.....98.798%
TOTAL	.....100.000%

### THIS PRODUCT CONTAINS:

13.50 lb 2,4-dichlorophenoxyacetic acid equivalent per ton (CAS 1928-43-4)

1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton (CAS 1918-00-9)

2.42 lb dithiopyr per ton (CAS 97886-45-8)

6.92 lb quinclorac per ton (CAS 84087-01-4)

Isomer Specific By AOAC Methods.

EPA Reg. No. 2217-927-32802

EPA Est. No. 32802-MO-1

DIMENSION® is a trademark of Dow AgroSciences.

Distributed by:

**Howard Johnson's Enterprises, Inc.**

700 W. Virginia Street, Milwaukee, WI 53204

Net Weight 50Lbs (22.68 kg)

## KEEP OUT OF REACH OF CHILDREN CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks, and
- chemical-resistant gloves.

#### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>.

### GUARANTEED ANALYSIS

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### General Precautions and Restrictions:

•Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled.

•Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.

•Cuatro with Dimension® plus Quinclorac is not sold, distributed, or registered within the state of New York. Do not use in the state of New York.

#### Product Description

Cuatro with Dimension® plus Quinclorac is formulated to feed your lawn and control most annual grasses and broadleaf weeds commonly found in home lawns. The product contains a lawn fertilizer and four different herbicides.

•The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake.

•The four herbicides combine to kill many different kinds of weeds at different stages of development.

•Grass weeds: Preemergent control of crabgrass, foxtails, and other annual grass weeds. Postemergent control of crabgrass.

•Broadleaf weeds: Preemergent control of chickweed, henbit, knotweed, shepherds-purse, spurge, woodsorel, and other broadleaf weeds. Postemergent control of broadleaf weeds including dandelion, chickweed, thistle and clover. A partial list of broadleaf weeds is presented on this label.

#### Where To Use

Cuatro with Dimension® plus Quinclorac may be used on the following established turf species:

Established Cool Season Turfgrass: Kentucky bluegrass, rough bluegrass (Poa trivialis), perennial ryegrass, fine fescues, tall fescue, seashore paspalum

Established Warm Season Turfgrass: Bermudagrass, buffalograss, zoysia-grass

Cuatro with Dimension® plus Quinclorac may NOT be used on: annual bluegrass (Poa annua), bahiagrass, bentgrass, carpetgrass, centipedegrass, dichondra, St. Augustine. Do not use this product on turfgrass where desirable clovers are present.

Cuatro with Dimension® plus Quinclorac may be used on warm-season and cool-season turfgrass in 4 use sites.

•**Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

•**Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.

•**Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs only), and office buildings.

•**Non-cropland sites** include highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations.

**Guidelines to follow:** The use of Cuatro with Dimension® plus Quinclorac on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

### WEEDS CONTROLLED

When applied as directed, this product will control and/or suppress\* the following weeds:

**Annual Grass Weeds:** Annual bluegrass (Poa annua), Annual ryegrass Barnyardgrass\*, Crabgrass (smooth, southern, large), Foxtail (green, yellow, giant), Goosegrass\* (crowfootgrass),

**A partial listing of established broadleaf weeds:** Bedstraw, Beggarticks, Black medic, Broadleaf plantain, Buckhorn plantain, Burclover, Burdock, Carpetweed, Chickweed, (common), Chickweed, (mouseear), Chicory, Clover, hop, Clover, strawberry Clover, (white), Curly dock, Daisy, oxeye, Dandelion, Dichondra, False dandelion, Ground ivy, Healall, Henbit, Knotweed, Lambsquarters, Lespedeza, Mallow, Matchweed, Morningglory, Mustard, Narrowleaf plantain, Oxalis, Parsley-piert, Pearlwort, Pennycress, Pepperweed, Pigweed, Plantain, Poison ivy, Poison oak, Pokeweed, Purslane (common), Ragweed, Red sorrel, Redroot pigweed, Sheep sorrel, Shepherds-purse, Slender plantain, Smooth dock, Smooth pigweed, Spurge, Thistle, Woodsorel, Yarrow and other broadleaf weeds

\* for acceptable control of barnyardgrass and goosegrass, two applications made at a 6 to 10 week interval are needed

### FIRST AID

**IF IN EYES**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Additional information in case of emergency, call toll free 1-877-800-5556

### Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

These chemicals (2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### When To Apply

This product may be applied as a single application or as split applications in the spring, early summer, or fall.

For annual grasses such as crabgrass (smooth, southern and large) and foxtail (green, yellow, and giant): Apply in the spring and early summer.

For best results, apply this product from 2 weeks before weed germination until the weed is 4 inches tall. For dense populations, make a second application no sooner than 30 days after the first application.

For broadleaf weed control: Spring applications: for chickweed, henbit, knotweed, shepherds-purse, spurge, woodsorel, and other broadleaf weeds. Fall applications: for chickweed, henbit, knotweed, shepherds-purse, spurge, woodsorel, and other broadleaf weeds. For best results, apply when broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F).

### How To Apply

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain.
- Poor results will occur if applied to dry lawns. When dry conditions exist, irrigate [or water] the turfgrass 24 hours before application.
- For best results, irrigate after the application. However do not irrigate [or water] sooner than 24 hours after application.
- Apply this product when the wind is calm.
- Do not spread granules onto flowers, vegetables, and ornamental shrubbery.
- Turf species listed on this label can be seeded, sodded, sprigged, or plugged into the treated areas 4 weeks after the application of this product.
- When treating newly planted areas, wait approximately 6 weeks after seedling emergence before applying this product.
- Delay mowing 2 days before and until 3 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area.

Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

### How Much To Use

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

Apply 2.4 to 4.5 pounds of product per 1,000 sq.ft. per application to established turfgrass.

The maximum broadcast application rate is 4.5 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum annual (seasonal) rate is 9.0 pounds of product per 1,000 sq.ft.

### SUGGESTED SPREADER SETTINGS

•For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.

Spreader setting to apply at the listed rate (lbs per 1000 sq ft)	Cone setting			
	2.4 lbs	3.6 lbs	4.5 lbs	
Accu-Pro 2000	6	K	M	N
SR2000	6	K	M	O
Lesco 021820	0.25 Closed	D	E	F
Spyker 76-22	4.3	4.2	4.4	4.6
PSB Bigfoot	A.5	K	K	M
Lely HR	II-B	5	5.75	7.75
Vicon 203		14	16	20

The above spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep in original container in a cool, dry area, away from extreme heat and cold.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. To avoid wastes, use the entire contents in this bag by application according to the label instructions. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container (bag). Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### LIMITED WARRANTY AND DISCLAIMER

**FOR USE ONLY AS DIRECTED.** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.