

Howard Johnson's

PROFESSIONAL CRABGRASS CONTROL

A GRANULAR FERTILIZER

CONTAINING TEAM* PRO HERBICIDE



FOR THE PREEMERGENCE CONTROL OF CRABGRASS, FOXTAIL, POA ANNUA, GOOSEGRASS & OTHER ANNUAL GRASS & BROADLEAF WEEDS IN ESTABLISHED TURF & LANDSCAPE ORNAMENTALS

ACTIVE INGREDIENTS:

benefin: N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine 0.43%

trifluralin: a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 0.43%

INERT INGREDIENTS: 99.14%

TOTAL: 100.00%

Contains 0.43 pounds of active ingredient per 50 pound bag

EPA Reg. No. 62719-289-32802

EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)

First letter of code printed across top of bag shows manufacturing establishment
*TEAM and TEAM logo are trademarks of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING:

Causes substantial but temporary eye injury. Harmful if absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency treatment information.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Personal Protective Equipment(PPE):

All loaders, applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves(such as or made of any waterproof material), and Goggles, face shield or safety glasses. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Users should: Wash hands after handling and before eating, drinking, chewing gum, using tobacco or using the toilet; Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing; Remove PPE immediately after handling this product.. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow people or pets to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated area (except those involved in the watering) until the watering-in is complete and the surface is dry.

GENERAL INFORMATION: This fertilizer product contains Team® Pro, a preemergence herbicide for control of annual grasses and broadleaf weeds in established turfgrasses and established landscape ornamentals.

WEEDS CONTROLLED:

Annual grasses:
annual bluegrass (*Poa annua*)
crabgrass, smooth (*Digitaria ischaemum*)
crabgrass, large or hairy (*Digitaria sanguinalis*)
green foxtail (*Setaria viridis*)
yellow foxtail (*Setaria glauca*)
barnyardgrass (watergrass) (*Echinochloa crus-galli*)
goosegrass (silver crabgrass or crowfootgrass) (*Elyusine indica*)

Broadleaf weeds:
chickweed, common (*Stellaria media*)
knotweed, prostrate (*Polygonum aviculare*)
lambquarters, common (*Chenopodium album*)
pigweed (*Amaranthus spp.*)
purslane, common (*Portulaca oleracea*)

WEEDS SUPPRESSED:

panicum, Texas (*Panicum texanum*)
prostrate spurge (*Euphorbia humistrata*)
creeping spurge (*Euphorbia serpens*)
woodsorrel (*Oxalis spp.*)
pusley, Florida (*Richardia scabra*)
spotted spurge (*Euphorbia maculata*)
creeping woodsorrel (*Oxalis corniculata*)

GUARANTEED ANALYSIS

Distributed By:
Howard Johnson's Enterprises, Inc.
700 W. Virginia St., No. 222
Milwaukee, WI 53204

USES:

ESTABLISHED TURFGRASS: When to apply: For optimum results, this product must be applied 1 to 2 weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons or for lawn care operator programs.

Application Timing	Application Rate	Specific Use Directions
Spring: To control spring-to-summer germinating annual weeds	174 to 349 lb / acre (4.00 to 8.01 lb / 1000 sq ft)	In cool season & warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination
Fall: To control fall-germinating weeds Split Application: For extended control of spring and summer annual weeds	174 lb / acre followed by 174 lb / acre (4.00 followed by 4.00 lb / 1000 sq ft) 233 lb / acre followed by 233 lb / acre (5.34 followed by 5.34 lb / 1000 sq ft)	In cool season turfgrass area & northern half of warm season turfgrass area, apply first application in spring 1 to 2 weeks before expected time of weed germination and make a second application 8 to 10 weeks later In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later

Amount Applied per acre (Pounds per 1000 square feet)	Minimum Time Following Last Application Before Reseeding
Single application of 174 lb / acre	8 Weeks
Single application of 349 lb / acre	12 to 16 Weeks
Split application of 174 lb / acre	8 to 10 Weeks
Split application of 233 lb / acre	10 to 12 Weeks
Season of Application	Pounds per 1000 sq ft
Spring application (Summer annuals)	4.00 to 8.01 lb / 1000 sq ft
Fall application (Winter annuals)	4.00 to 8.01 lb / 1000 sq ft

How to apply: Apply this product evenly over the turfgrass area using a properly calibrated drop-type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More

uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Activation: Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

IMPORTANT USE PRECAUTIONS: Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass. Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application. This product will not harm most nearby ornamentals. This product may be applied to the following established turfgrass species: Established cool season turfgrasses: Bentgrass fairways, Kentucky bluegrass, Perennial ryegrass, fescues. Established warm season turfgrasses: Bahiagrass, Bermudagrass, Centipedegrass, Fescue, tall, St. Augustinegrass, Zoysiagrass.

Broadcast Application Rates: Reseeding precautions: Team herbicide, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings, if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding, following the last application, should not be attempted until the applicable minimum time delay in the table above has been met.

ESTABLISHED ORNAMENTAL PLANTS IN LANDSCAPE SETTINGS: This product is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers. Application timing: Apply 1 to 2 weeks prior to the expected germination time of target annual weeds or to clean cultivated weed free areas. Do not apply this product to plantings when the foliage is wet. Soil preparation: This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate and hoe to remove existing weeds before applying this product. Application equipment and application techniques: See General Information section. Activation: Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application. Watering-in must be performed by the commercial applicator or the commercial applicator must provide the following instructions to the resident or owner in writing: The product must be watered in as soon as possible. Do not enter or allow other(including children or pets) to enter the treated areas(except those involved in the watering) until the watering-in is complete and the surface is dry.

Broadcast Application Rates:

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10-12 weeks following the initial application. This product may be used for annual weed control in landscape setting on the following established ornamental plant species:

Ornamental shrubs: andromeda, Japanese (*Pieris japonica*); arborvitae, American (*Thuja occidentalis*); azalea (*Rhododendron spp.*); barberry, Japanese (*Berberis thunbergii*); barberry, mento (*Berberis mentorensis*); boxwood, common (*Buxus sempervirens*); boxwood, littleleaf (*Buxus microphylla*); camellia, Japanese (*Camellia japonica*); camellia, sasanqua (*Camellia sasanqua*); cherrylaurel, American (*Prunus caroliniana*); cinquefoil (*Potentilla spp.*); cleyera, Japanese (*Cleyera japonica*); cotoneaster, cranberry (*Cotoneaster aciculata*); cotoneaster, zabel (*Cotoneaster zabelii*); deutzia (*Deutzia spp.*); elaeagnus, silverberry (*Elaeagnus pungens*); euonymus, spreading (*Euonymus kiautschovica*); euonymus, winged (*Euonymus alatus*); euonymus, wintercreeper (*Euonymus fortunei*); firethorn (*Pyracantha spp.*); forsythia (*Forsythia spp.*); guava, pineapple (*Feijoa sellowiana*); holly (*Ilex spp.*); honeysuckle (*Lonicera spp.*); indiahawthorn (*Ralphiolepis indica*); juniper (*Juniperus spp.*); laurel, mountain (*Kalmia latifolia*); lilac, common (*Syringa vulgaris*); mockorange (*Philadelphus spp.*); pittosporum, Japanese (*Pittosporum tobira*); privet (*Ligustrum spp.*); redcedar, eastern (*Juniperus virginiana*); rhododendron (*Rhododendron spp.*); spirea, vanhoutte (*Spiraea vanhouttei*); viburnum (*Viburnum spp.*); weigela (*Weigela spp.*); willow (*Salix spp.*); yew, anglojap (*Taxus media*); yew, Japanese (*Taxus cuspidata*); yewpine (*Podocarpus macrophyllus*).

Ornamental groundcover plantings: aaronsbeard (*Hypericum calycinum*); bellflower, adriatic (*Campanula elatines*); bellflower, poscharsky (*Campanula poscharskyana*); ceanothus (*Ceanothus spp.*); coreopsis (*Coreopsis spp.*); cotoneaster (*Cotoneaster spp.*); coyote brush (*Baccharis pilularis*); crown vetch (*Coronilla vana*); daisy trailing African (*Osteospermum fruticosum*); fern, asparagus (*Asparagus densiflorus*); gazania (*Gazania spp.*); germander (*Teucrium chamaedrys*); ice plant, largeleaf (*Carpobrotus edulis*); ivy, Algerian (*Hedera canariensis*); ivy, English (*Hedera helix*); lily-of-the-nile (*Agapanthus spp.*); lilyturf, bigblue (*Liriope muscari*); marigold (*Tagetes spp.*); myoporum (*Myoporum laetum*); plumbago, dwarf (*Ceratostigma umbaginoides*); rockrose (*Cistus spp.*); rosemary (*Rosmarinus officinalis*); rupterwort (*Herniaria glabra*); snow-in-summer (*Cerastium tomentosum*); speedwell (*Veronica spp.*); st. johnswort (*Hypericum coris*); stonecrop (*Sedum spp.*); strawberry, beach (*Fragaria chiloensis*); thrift (*Armeria maritima*); verbena (*Verbena spp.*); wirevine, creeping (*Muehlenbeckia axillaris*); yarrow, wolly (*Achillea tomentosa*).
Ornamental trees: almond (*Prunus dulcis*); apple, crabapple (*Malus spp.*); apricot (*Prunus armeniaca*); ash, white (*Fraxinus americana*); baldcypress (*Taxodium distichum*); birch, European white (*Betula pendula*); blackgum (*Nyssa sylvatica*); cherry (*Prunus spp.*); chestnut, Chinese (*Castanea mollissima*); cottonwood (*Populus deltoides*); dogwood, flowering (*Cornus florida*); dogwood, kousa (*Cornus kousa*); douglasfir (*Pseudotsuga menziesii*); fir, balsam (*Abies balsamea*); hemlock, Canada (*Tsuga canadensis*); honeylocust (*Gleditsia triacanthos*); larch, Japanese (*Larix kaempferi*); locust, black (*Robinia pseudoacacia*); maple, Norway (*Acer platanoides*); maple, red (*Acer rubrum*); maple, silver (*Acer saccharinum*); maple, sugar (*Acer saccharum*); oak, pin (*Quercus palustris*); oak, scarlet (*Quercus coccinea*); peach (*Prunus persica*); pine, Austrian (*Pinus nigra*); pine, eastern white (*Pinus strobus*); pine, Japanese black (*Pinus thunbergiana*); pine, loblolly (*Pinus taeda*); pine, red (*Pinus resinosa*); pine, scotch (*Pinus sylvestris*); planetree, London (*Platanus acerifolia*); plum (*Prunus spp.*); redbud, eastern (*Cercis canadensis*); spruce, Colorado (*Picea pungens*); spruce, Norway (*Picea abies*); sycamore (*Platanus occidentalis*); tuliptree (*Liriodendron tulipifera*); walnut, black (*Juglans nigra*).
Roses and other established flowers: African daisy; aster (perennial); balsam; blackeyed susan; calendula; carnation; centaurea, velvet; crysanthemum; coreopsis; cornflower; cosmos; dahlia; dianthus; dusty miller; floss flower; forget-me-not; four o'clock; gaillardia; gladiolus; golden glow; inpatiens; ixora; lobelia; lupine; marigold; marigold, cape; morningglory; nasturtium; nicotiana; petunia; phlox; pincushion flower; poppy, California; portulaca; rose; salvia; shasta daisy; snapdragon; snow-on-the-mountain; stock; sunflower; sweet alyssum; sweet pea; sweet sultan; sweet william; vinca; yarrow; zinnia.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, contain material & dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

SPREADER SETTINGS:

To apply at listed rate:

The settings shown here are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed, and equipment condition.

Calibrate your spreader before use to ensure the correct application rate.

Spreader setting to apply at the listed rate (lbs per 1000 sq ft)	Spreader	Cone setting	4 lbs	5.34 lbs	8 lbs
Accu-Pro 2000	6	M 1/2	N 1/2	Q	
SR2000	6	N	O 1/2	Q 1/2	
Lesco 021820	0.25 Closed	G	F 1/2	I	
Spyker 76-22		4.8	4.7	5.3	
PSB Bigfoot	A.5	M 1/2	L 1/2	N	
Lely HR	II-B	6.5	6.5	8	
Vicon 203		17	19	25	

Terms & Conditions of Use: If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return the unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other use constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies. **Warranty Disclaimer:** Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent consistent with applicable law Seller makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty. **Inherent Risks of Use:** It is impossible to eliminate all risks associated with the use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornados, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer. **Limitation of Remedies:** To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following: (1) Refund of purchase price paid by buyer or user for product bought, or, (2) Replacement of amount of product used. Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses. The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.