

Howard Johnson's SEED SAFE TURF CARE

TUPERSAN® PRE-EMERGENCE HERBICIDE PLUS TURF FOOD

MAY BE USED ON NEWLY SEEDED LAWNS

When used as directed, will not harm desirable lawn grass seed. Yet kills annual weed grasses such as Crabgrass, Foxtail & Barnyardgrass (Watergrass) on home lawns, golf courses, ornamental & recreational turfgrass.

This herbicide is not for use on Bermudagrass or Bentgrass.

ACTIVE INGREDIENT:
Siduron; 1-(2-methylcyclohexyl)-3-phenylurea . . . 3.71%
INERT INGREDIENTS: 96.29%
TOTAL: 100.00%

EPA Reg. No. 32802-28
EPA Est. No. 32802-WI-1 (A), 32802-MO-1 (B)
First letter of bag code shows establishment where manufactured.
TUPERSAN® is a registered trademark of Gowan Company

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs. Do not use on turf to be used for feed or food purposes.
ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

GUARANTEED ANALYSIS

First Aid	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Hot line number:	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

Manufactured & Guaranteed by:

Howard Johnson's Enterprises, Inc.

700 W. Virginia Street, Milwaukee, WI 53204

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its label. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This pre-emergence herbicide plus turf food may be used on newly seeded lawns. When used as directed, will not harm desirable lawn grass seed - yet kills annual weed grasses on home lawns, golf courses, parks, roadsides, athletic fields, all ornamental and recreational turfgrass.

WEEDS CONTROLLED: Annual weed grasses such as Crabgrass (smooth and hairy), Foxtail (green and yellow) Barnyardgrass (Watergrass). This product will not control *Poa annua*. This product will not control established weeds that are already growing.

WHERE TO USE: For use on Bluegrass, Fescue, Redtop, Perennial Ryegrass, Smooth Brome, Orchardgrass and certain Bentgrass strains (Penncross, Seaside, Highland, Astoria, Nimisila, C1, C7 and C19). Do not use on other Bentgrass varieties since injury may occur. Do not use on Bermudagrass.

HOW MUCH AND WHEN TO USE:

FOR NEW SPRING SEEDINGS: Apply 2.5 pounds per 1000 square feet to treat and feed lawn where bare ground or very thin turf is seeded in the spring and 2 weeks prior to weed seed germination. Follow good cultural practices of keeping area moist to help grass seed germinate and also to activate the pre-emergent herbicide and to get maximum benefit from the lawn food.

FOR NEW "FALL" SEEDING OR SEEDING TO ESTABLISHED LAWNS IN THE SPRING: Apply 5 pounds per 1000 square feet to treat and feed turf. Apply in spring prior to the expected emergence of the annual weed grasses. This will vary from early February in the south to early March in the north. Follow the application with a thorough watering (1/2 inch) to activate the herbicide.

HOW TO APPLY: To insure best weed control and proper fertilization the lawns should be free of leaves, grass clippings or thatch which might interfere with the pre-emergence action of the herbicide. Complete all cultural practices such as aeration before application. Apply with a properly calibrated fertilizer spreader. If it does not rain within 3 days after application, water the area thoroughly to get desired results.

RESEEDING TREATED AREAS: This product may be applied to the soil at the same time as sowing grass seed. When used as directed on listed turfgrasses this product will not affect the germination of those seedling grasses.

SPREADER SETTINGS: Make sure your spreader is operating correctly. Spreader settings are for use with new equipment. Since proper application is based on the spreader's condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require adjustment. To check your spreader to apply 5 pounds per 1000 square

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

For optimum pre-emergence weed control apply at the appropriate time

Jan Feb Mar Apr May June July Aug Sept Oct Nov Dec

1. [REDACTED]

2. [REDACTED]

- For crabgrass, Foxtail & other grassy weeds in NORTHERN STATES
- For crabgrass, Foxtail & other grassy weeds in SOUTHERN STATES

SPREADER SETTINGS:

The settings shown here are approximate for new equipment. Your spreader may need calibration, due to differences in terrain, walking speed, and equipment condition.

Calibrate spreader to ensure correct application rate. Clean and lubricate your spreader after each use.

SPREADER SETTINGS			
Spreader	Cone Setting	2.5 lbs	5 lbs
Accu-Pro	6	L 1/2	N 1/2
SR2000	6	L 1/2	O 1/2
Lesco	0.25 closed	D 1/2	E
Spyker 76-22		4.3	4.7
PSB Bigfoot	A.5	J 1/2	L 1/2
Lely HR	II-B	5.25	6.5
Vicon		15	19

feet, place 2.5 pounds of this product in your spreader, using setting shown, and apply to 500 square feet. If you run out of granules before you cover the pre-measured areas, close the spreader setting by 1/2 number. If you have granules left over, open (increase) the spreader setting by 1/2 number. For further help refer to your spreader guide.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

02822

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container in a cool, dry, locked place out of reach of children.

Pesticide disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.