



Howard Johnson's

CRABGRASS CONTROL PLUS WITH 0.37% BARRICADE®

FOR SELECTIVE PREEMERGENCE CONTROL OF GRASS & BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF GRASSES (EXCLUDING GOLF COURSE PUTTING GREENS), LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS & WILDFLOWER PLANTINGS. STOPS MANY UGLY WEED GRASSES BEFORE THEY GROW. PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL, & MORE.

ACTIVE INGREDIENT:

Prodiamine: [N³,N³-Di-*n*-propyl-2,4-dinitro-6-(trifluoromethyl)-*m*-phenylenediamine]0.37%

OTHER INGREDIENTS:99.63%
TOTAL:100.00%

EPA Reg. No. 32802-66

EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (C),
8378-IN-1 (D), 9404-FL-1 (E)

First letter of bag code shows establishment where manufactured.

BARRICADE® is a registered trademark of Syngenta Crop Protection, Inc.

Manufactured for: Howard Johnson's Enterprises, Inc.
9675 S 60th Street, Franklin, WI 53132

Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

Agricultural Chemical: Do not ship or store with food, feed, drugs or clothing. For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424-9300.

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS: This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

| | <u>First Aid</u> |
|--|--|
| If in eyes | <ul style="list-style-type: none"> * Hold eye open and rinse slowly and gently with water for 15-20 minutes. * Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. * Call a poison control center or doctor for treatment advice. |
| In on skin or clothing | <ul style="list-style-type: none"> * Take off contaminated clothing. * Rinse skin immediately with plenty of water for 15-20 minutes. * Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> * Call a poison control center or physician immediately for treatment advice. * Have a person sip a glass of water if able to swallow. * Do not induce vomiting unless told to do so by the poison control center or physician. * Do not give anything by mouth to an unconscious person. |
| If inhaled | <ul style="list-style-type: none"> * Move person to fresh air. * If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. * Call a poison control center or doctor for further treatment advice. |
| | <u>Hotline number</u> |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378. | |

GUARANTEED ANALYSIS

25-0-5

Total Nitrogen (N) **25%**

25.00% Urea Nitrogen

Soluble Potash (K₂O) **5%**

Derived From: Muriate of Potash,
Urea

Chlorine (Cl), not more than **6.00%**

F315

Bulk Denisty
59

Item Number
FGB55648
100206793

Net Weight 50 lbs (22.70 kg)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit²; itchgrass; johnsongrass (from seed); junclegrass; knothead; Kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³; shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsrrel, yellow (from seed).

¹ In many areas where not prohibited by fertilizer use restrictions** a single application of 4.1 to 9.3 pounds / 1000 square feet will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4.1 to 6.2 pounds / 1000 square feet, to areas not prohibited by fertilizer use restrictions, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see Minimum and Maximum Application Rate table).

² For this weed, apply in late summer, fall or winter prior to germination.

³ Suppression only.

**Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days after application.

In the State of Florida, this product should be applied at rates not exceeding those recommended in the document titled Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil (not thatch plus soil).

| Spreader setting to apply at the listed rate (lbs per 1000 sq ft) | Cone setting | 3.0 lbs | 4.0 lbs | 4.6 lbs | 6.2 lbs | 9.3lbs |
|---|--------------|---------|---------|---------|---------|--------|
| | | 6 | M 1/2 | N 1/2 | O 1/2 | Q 1/2 |
| Accu-Pro 2000 | 6 | L 1/2 | N | O 1/2 | P | R |
| SR2000 | 6 | L 1/2 | N | O 1/2 | P | R |
| Lesco 021820 | 0.25 Closed | D 1/2 | E 1/2 | F 1/2 | G 1/2 | Q 1/2 |
| Spyker 76-22 | | 4.3 | 4.5 | 4.7 | 4.9 | 5.5 |
| PSB Bigfoot | A.5 | J 1/2 | K 1/2 | L 1/2 | M | N 1/2 |
| Lely HR | II-B | 5.25 | 6 | 6.5 | 7 | 8.5 |
| Vicon 203 | | 15 | 17 | 19 | 21 | 27 |

| Turf Species | Minimum and Maximum Application Rates (by Turf Grass Species) ¹ | |
|---|--|--------------------------------------|
| | Minimum Rate | Maximum Annual Rate ² |
| Bahiagrass ³ , Centipedegrass ³ Bermudagrass ³ , Zoysiagrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) | 2.7 pounds per 1000 sq. ft. | 9.3 pounds per 1000 sq. ft. per year |
| Buffalograss Kentucky Bluegrass Perennial Ryegrass | 2.7 pounds per 1000 sq. ft. | 6.2 pounds per 1000 sq. ft. per year |
| Creeping Red Fescue | 2.7 pounds per 1000 sq. ft. | 4.6 pounds per 1000 sq. ft. per year |
| Creeping Bentgrass | 2.7 pounds per 1000 sq. ft. | 4.0 pounds per 1000 sq. ft. per year |

¹ These are the maximum rates per calendar year.
² In Florida: Do not exceed 2 applications per year to Bahiagrass and Centipedegrass in the north. do not exceed 3 applications per year to Centipedegrass in the central and south regions. Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.
³ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 lbs / 1000 square feet. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

| Use Rate lbs / 1000 sq ft | Months Before Overseeding | | |
|------------------------------|---------------------------|------------|-------|
| | North | Transition | South |
| 2.7 lbs | 4 | 4 | 4 |
| 3 lbs | 4 | 4 | 4 |
| 4 lbs | 5 | 4 | 4 |
| 4.6 lbs | 6 | 5 | 5 |
| 4.9 lbs | --- | 6 | 6 |
| 6.2 lbs | --- | 7 | 7 |
| 7 lbs | --- | --- | 9 |
| 8 lbs | --- | --- | 10 |
| 9.3 lbs | --- | --- | 12 |

Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 lbs / 1000 square feet per year.

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds, and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

NOTICE: This product contains chemical known to the State of California to cause cancer and birth defects and other reproductive harm.

| SCIENTIFIC NAME | COMMON NAME | SCIENTIFIC NAME | COMMON NAME | SCIENTIFIC NAME | COMMON NAME |
|---------------------------|---------------------------------|--------------------------|-----------------------------|---------------------------|---|
| Abies spp | Fir species** (Balsam, etc.) | Hedera helix | English Ivy | P. tobira | Japanese Pittosporum |
| Acer palmatum | Japanese Maple | Hibiscus rosa-sinensis | Chinese Hibiscus | Podocarpus macrophyllus | Japanese Yew |
| A. platanoides | Norway Maple | Ilex cornuta | Chinese Holly | Prunus laurocerasus | English Laurel |
| Actinidia chinensis | Kiwi* | I. crenata | Japanese Holly | Prunus sp. | Almond, Apricot, Nectarine Peach, Plum and Prune* |
| Agapanthus africanus | Lily-of-the-Nile (African Lily) | I. opaca | American Holly | | |
| Arctostaphylos densiflora | Vine Hill Manzanita | Iris spp. | Iris species** | Pyraecanthia coccinea | Firethorn (Scarlet) |
| Arctotheca calendula | Cape Weed | Jasminium nudiflorum | Winter Jasmine | P. fortuneana | Firethorn |
| Aucuba japonica | Japanese Aucuba | Juniperus chinensis | Chinese Juniper | P. koidzumii | Firethorn |
| Berberis gladwynensis | Barberry | J. conferta | Shore Juniper | Pyrus sp. | Bradford Pear sp. |
| B. julianae | Wintergreen Barberry | J. horizontalis | Creeping Juniper | Quercus spp. | Oak species |
| B. mentorensis | Mentor Barberry | Juglans sp. | Walnut | Rapiclopsis indica | Indian Hawthorne |
| B. thunbergii | Japanese Barberry | Justicia brandegeana | Shrimp Plant | Rhododendron | 'Coral Bells' |
| Buxus microphylla | Japanese Boxwood | Lagerstromia indica | Crape Myrtle | (including Azalea) | 'Formosa,' 'Hino-crimson,' 'PJM,' 'Roseum Elegans' |
| Callistemon viminalis | Weeping Bottlebrush | Ligustrum amurense | Arner Privet | Rosa banksiae | Lady Bank's Rose |
| Calluna vulgaris | Scotch Heather | L. lucidum | Japanese Privet | Rosmarinus officinalis | Rosemary |
| Carpobrotus edulis | Hottentot Fig (Ice Plant) | L. luridum | Glossy Privet (Waxleaf) | Santolina viensis | Santolina |
| Cassia artemisoides | Feathery Cassia | L. muscari | Big Blue Lily, Silver Mound | Sedum album | Stonecrop |
| Ceanothus rigidus | Wild Lilac | L. japonica | Japanese Honeysuckle | Syzygium paniculatum | Japanese Brushcherry |
| Chamaecyparis pisifera | False Cypress | L. tatarica | Tatarian Honeysuckle | Taxus cuspidata | Japanese Yew |
| Cleyera | Cleyera | L. maleophora luteola | Ice Plant | T. media | Yew |
| Citrus spp. | Citrus species* | Malus sp. | Crabapple* | Thuja occidentalis | American Arborvitae |
| Cornus florida | Flowering Dogwood | Nandina domestica | Heavenly Bamboo | Trachelospermum asiaticum | Star Jasmine |
| C. stolonifera | American Dogwood | Olea europaea | Olive* | Tsuga canadensis | Canada Hemlock |
| Cortaderia selloana | Pampas Grass | Ophiopogon japonicus | Mondo Grass | Viburnum japonicum | Japanese Viburnum |
| Cotoneaster apiculatus | Cranberry Cotoneaster | Osteospermum fruticosum | Trailing African Daisy | V. odoratissimum | Sweet Viburnum |
| C. buxifolius | Cotoneaster | Oxydendrum arboreum | Sourwood | V. plicatum | Japanese Snowball |
| C. dammeri | Seaberry Cotoneaster | Persea americana | Avocado* | V. rufidum | Canary Island Viburnum |
| C. microphyllus | Rockspray Cotoneaster | Photinia fraseri | Fraser's Photinia (Redtip) | V. tinus** | Laurastinus |
| C. taegus spp. | Hawthorne | Picea spp. | Spruce species* | V. trilobium | Cranberry Bush |
| Cupressus sempervirens | Italian Cypress | Pieris japonica | Lily-of-the-Valley Bush | V. wrightii | Leatherleaf Viburnum |
| Delasperra alba | White Trailing Ice Plant | Pinus brutia | Calabrian Pine | Vinca major | Greater Periwinkle, Vinca |
| Delphinium spp. | Larkspur, Blue Elf | P. canariensis | Canary Island Pine | Vinca minor | Common Periwinkle, Vinca |
| Dodonea viscosa | Hop Bush | P. halepensis | Aleppo Pine | Vitis sp. | Grape* |
| Elaeagnus pungens | Silverberry | P. nigradiata | Austrian Pine | Weigela florida | Old Fashioned Weigela |
| Euonymus fortunei | Wintercreeper | P. thunbergiana | Japanese Black Pine | Yucca aloifolia | Spanish Bayonet, Yucca |
| E. japonica | Evergreen Euonymus | P. strobus | Spreading Euonymus | Y. filamentosa | Adam's Needle, Yucca |
| E. kiautschovica | Spreading Euonymus | P. sylvestris | Scotch Pine | | *Do not use on food producing trees, vines or plants. |
| Fatsia japonica | Japanese Aralia | Pistachio sp. | Pistachio* | | **Not for use on container grown plants. |
| Forsythia intermedia | Border Forsythia | Pittosporum rhombifolium | Queensland Pittosporum | | |
| F. viridissima | Greenstem Forsythia | | | | |
| Gardenia jasminoides | Gardenia, Cape-Jasmine | | | | |

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. If not available then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Warranty: The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc. the manufacturer, or the seller.
To the extent consistent with applicable law, the products sold to you are furnished "as is" by Howard Johnson's Enterprises, Inc. The manufacturer and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Howard Johnson's Enterprises, Inc., the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc. the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc. or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product. To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc. the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.