



# Howard Johnson's ECHELON™

## 0.3% Granular Herbicide on Fertilizer

For Preemergence and early Postemergence Control of Crabgrass, selected annual grasses, sedges, and annual broadleaf weeds in Turf Sites -Residential and Institutional Lawns, Athletic Fields, Commercial Sod Farms, Golf Course Fairways and Roughs, Roadsides, Utility right-of-ways, Railways and Industrial areas

**Active Ingredient:** Sulfentrazone-N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide . . . . . 0.1%  
 Prodiamine: [N3,N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine . . . . . 0.2%  
 Other Ingredients: . . . . . 99.7%  
 100.0%

Echelon 0.3% Granular contains 0.003 pounds of active ingredient per pound of product U.S. Patent No. 4,818,275

EPA Reg. No. 279-3322-32802  
 EPA EST.No. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (C)

FLORIDAAPPLICATIONS: For commercial application to urban turf or lawns in FL, we recommend you follow the Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries, available on the internet at [http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP\\_Bookfinal.pdf](http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Bookfinal.pdf)  
 FLORIDAAPPLICATIONS: Fro use on golf cours or other athletic turf in FL, we recommend you follow SL 191, "Recommendations for N, P, K & Mg for Golf Course and Athletic Field Fertilization Base on Mehlich I Extractant, available on the internet at <http://edis.ifas.ufl.edu/SS404>

Distributed for: **Howard Johnson's Enterprises, Inc.**  
 9675 S 60th Street, Franklin, WI 53132

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**FIRST AID**

IF INHALED	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF IN EYES	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>

**HOTLINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call 1-800-331-3148 for emergency medical treatment information.

**For Information Regarding the Use of this Product Call Howard Johnson's 1-800-298-4656**

## PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals) CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing dust. Do not get on skin, in eyes or on clothing. Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.  
**Personal Protective Equipment (PPE)**  
 Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: • Users should remove clothing immediately if product gets inside. Then wash thoroughly and put on clean clothing.
--

**Environmental Hazards**  
 This product is toxic to marine/estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.  
**Groundwater advisory:** Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.  
 Do not use on coarse soils classified as sand which have less than 1% organic matter.  
**Surface water advisory:** Sulfentrazone can contaminate surface water through spray drift. Under some conditions, sulfentrazone may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-lying tile drainage systems that drain to surface waters.

## GUARANTEED ANALYSIS 16-0-6

Total Nitrogen (N) 16%  
 16.00% Urea Nitrogen  
 Soluble Potash (K2O) 6%  
 Derived From: Muriate of Potash, Urea  
 Chlorine, (Cl), not more than 6.00%

F315  
 Bulk Density 66  
 Item Number FGB56561  
 100507480

## Net Weight 50 lbs (22.68 kg)

**DIRECTIONS FOR USE**  
 It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**Agricultural Use Requirements**  
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. These requirements only apply to uses of this product that are covered by the Worker Protection Standard.  
 Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.  
 PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

**Non-Agricultural Use Requirements**  
 The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.  
 Re-entry Statement: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Storage:** Store product in original container only, away from other pesticides, fertilizer, food or feed. Reseal open bag by folding top down and securing. Store in a cool, dry place and avoid excess heat.  
**Spill:** In case of spill, avoid contact, isolate area and keep out animals and unprotected persons.  
**Confine spills. Call FMC: (800) 331-3148.**  
 To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.  
**Pesticide Disposal:** If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance.  
**Container Disposal:** Paper and Plastic Bags: Non-refillable. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill, or by incineration, or by burning. Do not burn unless allowed by state or local authorities. If burned, stay out of smoke.

**GENERAL INFORMATION**  
**Turf Use Instructions**  
 Howard Johnson's Echelon is a selective pre-emergence and post-emergence herbicide for control of crabgrass, annual grass and broadleaf weeds and sedges in turf sites including residential and institutional lawns, athletic fields, commercial sod farms, golf course fairways and roughs, roadsides, utility right-of-ways, railways and industrial areas.

Howard Johnson's Echelon is a granular formulation containing 0.003 lbs. of active ingredient per pound of formulated product. The mode of action of Howard Johnson's Echelon herbicide involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings.

**SPECIFIC USE INSTRUCTIONS IN TURFGRASS**  
**Established Turf**  
 This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.  
 Broadcast Howard Johnson's Echelon with suitable application equipment to ensure uniform coverage over the treatment area. A drop or rotary spreader designed to apply granular herbicide or insecticide is the preferred application of this product. Calibrate application equipment prior to application. Consult manufacturer operation manual for proper calibration and operation of application equipment. Apply 2.87 to 8.60 pounds per 1,000 square feet to control broadleaf weeds and grasses. See Table 1 for application rates based on turf grasses.  
**When applied as directed under the conditions described, the following established grasses are tolerant to Howard Johnson's Echelon herbicide at the recommended use rates in a range from 0.250 to 1.125 lb. ai/acre (83 to 375 pounds/acre or 1.91 to 8.6 pounds/1,000 sq. ft.). Do not exceed 1.125 lb. ai/acre (375/acre) per calendar year.**

Table 1. Tolerant grasses

Grass Type	Recommended Use Rates Do not exceed maximum use rates per turf species in a single application		
	Lb. ai/acre	Lbs/ 1000 ft <sup>2</sup>	Lbs / acre
<b>Cool Season Grasses</b>			
Bentgrass, creeping <sup>1</sup>	0.25 - 0.375 <sup>3</sup>	1.90 - 2.87	83-125 lbs
Fescue, fine (Festuca rubra)	lb. ai.	lbs	
Ryegrass, perennial (Lolium perenne)			
Bluegrass, Kentucky (Poa pratensis)	0.57 - 0.75 <sup>3</sup>	4.36 - 5.74	190-250 lbs
Fescue, tall <sup>2</sup> (Festuca arundinacea)	lb. ai	lbs	
<b>Warm Season Grasses</b>			
Bahia grass (Paspalum notatum)			
Buffalograss (Buchloe dactyloides)			
Carpetgrass (Axonopus affinis)			
Centipedegrass (Eremochloa ophiuroides)	0.57 - 0.75 <sup>3</sup>	4.36 - 5.74	190-250 lbs
Kikuyugrass (Pennisetum clandestinum)	lb. ai.	lbs	
Seashore Paspalum (Paspalum vaginatum)			
Zoysiagrass (Zoysia japonica)			
Bermudagrass (Cynodon dactylon) & hybrids	0.75 - 1.125 <sup>3</sup> lb.ai.	5.74 - 8.6 lbs	250 - 375 lbs

- If bentgrass is mowed lower than 0.5 inches do not apply Howard Johnson's Echelon.
- Use of this product on certain varieties of Cheewing Fine Fescue or Tall Fescue cultivars may result in undesirable injury.
- Howard Johnson's Echelon may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate per turf type or 1.125 lbs. ai/acre (375 ounds/acre), whichever is lower.
  - \* Use of this product on certain cultivars of Cheewings Fescue, Fine Fescue or Tall Fescue ay result in undesirable injury.
  - \* Echelon application may cause temporary discoloration to exposed leaf surfaces on certain cultivar of zoysia grass. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply Echelon on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth.
  - \* Howard Johnson's Echelon has demonstrated tolerance on both cool and warm season turfgrasses. However not all varieties and/or cultivars have been evaluated. Turfgrass managers desiring to treat newly released varieties should first apply Echelon to a small area prior to treatment of larger areas.

Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	(Echinochloa crus-galli)	Lambsquarters,	(Chenopodium album)
Bluegrass annual <sup>1</sup>	(Poa annua),	common	
Carpetweed	(Mollugo verticillata)	Lovegrass	(Eragrostis spp.)
Carolina Geranium	Geranium carolinianum	Nutsedge, yellow	(Cyperus esculentus)
Chickweed, common <sup>2</sup>	(Stellaria media)	Panicum(Texas,Fall)	(Panicum spp.)
Chickweed,mousear	(Cerastium vulgatum)	Pigweed, Redroot	(Amaranthus retroflexus)
	(from seed)	Purslane, Common	(Portulaca oleracea)
Crabgrass	Digitaria spp.)	Pusley, Florida	(Richardia scabra L.)
(large, Smooth) <sup>3</sup>		Sedge, globe	(Cyperus globulosus)
Crowfootgrass	(Dactyloctenium aegyptium.)	Sedge, cylindrical	(Cyperus retrorsus)
Cupgrass, Woolly	(Eriochloa villosa)	Sedge, Surinam	(Cyperus surinamensis)
Foxtails, Annual	(Setaria spp..)	Sedge, Texas	(Cyperus polystachyos)
Goosegrass <sup>4</sup>	(Eleusine indica)	Shephardspurse <sup>2</sup>	(Capsella bursa-pastoris)
Hairy Bittercress	Cardamine hirsute,	Signalgrass, broadleaf	(Bracharia playphylla)
Henbit <sup>2</sup>	(Lamium amplexicaule )	Speedwell	(Veronica spp.)
Itchgrass	(Rottboellia exaltata)	Sprangletop	(Leptochola spp.)
Knotted, prostrate	(Polygonum aviculare)	Spurge, prostrate	(Euphorbia humistrata)
Kochia	(Kochia scoparia)	Witchgrass	(Panicum capillare L.)
Kyllinga, green	(Kyllinga brevifolia),	Woodsorrel, yellow	(Oxalis stricta)
Kyllinga, false	(Kyllinga gracillma)	(from Seed)	

- Poa annua is a winter annual. Applications of Howard Johnson's Echelon should be made in August or September to established, non overseeded turf before Poa Annua germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August and September are approximate timings.
- Apply Howard Johnson's Echelon in late summer, fall or winter before weed germinates for control.
- Crabgrass control in cool season grasses from fall application:  
 Areas where the ground is subject to winter freezes, Howard Johnson's Echelon may be applied in the fall at rates listed in Table 1 (83-250 pounds per acre) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring.  
 In areas of extended growing season and heavy pressure weed control will be enhanced by second application of Howard Johnson's Echelon 60-90 days after initial application
- In areas of extended growing season and heavy pressure weed control will be enhanced by second application of Howard Johnson's Echelon 60-90 days after initial application

**Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds and Sedges**  
 Howard Johnson's Echelon herbicide will control or suppress the weeds listed in Table 3 when applied shortly after weeds have emerged. Do not exceed the application rate specified for the turf species in Table 1.  
**Weeds Controlled or Suppressed When Application Made Postemergence to Weeds (Table 3)**

Common Name	Scientific Name	Controlled	Suppressed
Bittercress	(Cardamine spp.)		X
Black medic	(Medicago lupulina)		X
Buttercup	(Ranunculus spp.)		X
Carolina geranium	(Geranium carolinianum)	1	X
Carpetweed	(Mollugo verticillata)		X
Chickweed, common	(Stellaria media)	1	
Chickweed, mousear	(Cerastium vulgatum)		X
Cinquefoil	(Potentilla spp.)		X
Clover	(Trifolium spp.)		X
Crabgrass(large & Smooth)	(Digitaria spp.)	2	X
Cudweed	(Cynophallum spp.)		X
Dandelion	(Taraxacum officinale)		X
Dock, Curly	(Rumex crispus)		X
Evening primrose	(Oenothera biennis)		X
Fiddleneck	(Amsinckia spp.)		X
Filaree	(Erodium spp.)		X
Goldenrod	(Solidago spp.)	1	
Goosegrass	(Eleusine indica)	2	
Ground ivy	(Glechema hederacea)	1	
Henbit	(Lamium amplexicaule)		X
Knotted, prostrate	(Polygonum aviculare)	1	X
Kochia	(Kochia scoparia)	1	
Lambsquarters, common	(Chenopodium album)		X
Lawn burweed	(Solliva pterosperma)		X
Lespedeza, common	(Lespedeza striata)		X
Mallow, common	(Malva neglecta)		X
Nutsedge, Yellow	(Cyperus esculentus)	C	
Parsley piert	(Aichemilla arvensis)	1	
Pigweed, Redroot	(Amaranthus retroflexus)	1	
Pigweed, Tumble	(Amaranthus albus)	1	
Plantain, broadleaf	(Plantain matricarionoides)		X
Plantain, buckhorn	(Plantago lanceolata)	1	
Puncture weed	(Tribulus terrestris)		X
Purslane, common	(Portulaca oleracea)		X
Pusley, Florida	(Richardia scabra)	1	
Redweed	(Melochia corchorifolia)		X
Rocket, London	(Sisymbrium iro)		X
Smartweed, Pennsylvania	(Polygonum pensylvanicum)	1	
Sorrel, Red	(Rumex acetosella)		X
Speedwell	(Veronica spp.)	1	
Spurge, (annuals)	(Euphorbia spp.)	1	
Spurge, prostrate	(Euphorbia humistrata)	1	
Spurge, spotted	(Euphorbia maculata)	1	
Star of Bethlehem	(Ornithogalum umbellatum)	1	
Velvetleaf	(Abutilon theophrasti)		X
Violet, wild	(Viola pratincola)		X
Wild garlic	(Allium vineale)		X
Wild onion	(Allium canadense)		X
Woodsorrel, creeping	(Oxalis corniculata)		X
Woodsorrel, yellow	(Oxalis stricta)	1	

- Weeds are suppressed at lower rates (<250 lbs/acre), for optimum control apply rates of at least 250 lbs per acre in a single application Do not exceed the application rate specified for the turf species in Table 1
- Echelon controls crabgrass and goosegrass when applied postemergent to newly emerged weeds in the 1-4 leaf stage of development.

**Application to Reseeded, Overseeded or Sprigged areas**  
 Reseeding, overseeding or sprigging of treated areas within three to twelve (3-12) months after application of this product could inhibit the establishment of desirable grasses. [See table for exact intervals from application]. Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.  
**Overseeded Intervals after Howard Johnson's Echelon Applications (Table 4 )**

Total Amount of Echelon Applied	Interval (Months Before Overseeding)		
Pounds Product/Acre	North	Transition	South
83 - 125	3	3	3
166 - 250	4	4	4
375	6	5	5

**Sod Production**  
 It is recommended that sod be established for at least six (6) months before an application of Howard Johnson's Echelon herbicide.  
 Do not apply this product within three (3) months of harvest.  
**Other Use Precautions**  
 Do not apply to golf course putting greens or tees.  
 Do not use on turfgrasses other than those listed on this label.  
 Do not graze or feed livestock forage cut from areas treated with Howard Johnson's Echelon herbicide.  
 Do not apply directly to landscape ornamentals or ornamental beds.  
 Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species.

Do not apply to creeping bentgrass if mowing height is less than 0.5 inch.  
 Do not harvest treated sod before 90 days after application.  
 Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.  
 Do not apply with adjuvants or surfactants.

**ADDITIONAL USE INSTRUCTIONS**  
**Railroad Rights-of-Way**  
 Controls many weeds and maintains bare ground on railroad rights-of-way, including railroad yards, railroad crossings and railroad bridge abutments.  
**Highway, Roadside, Pipeline and Utility Rights-of-Way**  
 Controls many weeds and maintains bare ground in highway, roadside, pipeline and utility rights-of-way. Such areas would include, but are not limited to, guardrails, road shoulders, electric utility substations, pipeline pumping stations, around electric transmission towers, around distribution line poles plus other areas where complete vegetation control is desired.  
**Industrial Areas, Fence Rows and Other Non-crop Sites**  
 Controls weeds and maintains bare ground in industrial areas including production facilities, tank farms, storage areas, parking areas, lumber yards, airports, military installations, along fence rows, and in similar non-crop sites where complete vegetation control is needed.

**METHOD AND RATE OF APPLICATION FOR ADDITIONAL USES**  
 For residual control of germinating weeds in non-crop land, apply as a broadcast treatment using 250 to 375 pounds (0.75 to 1.125 pounds active ingredient) per acre by ground in a suitable granular applicator.  
 DO NOT apply Howard Johnson's Echelon Herbicide to soils classified as sand with less than 1% Organic Matter.

**Timing**  
 For best results, apply Howard Johnson's Echelon alone or in combination with other herbicides for residual control of weeds in late summer, fall, or early spring to insure adequate moisture for soil activation.  
**Weed Control**  
 Howard Johnson's Echelon when applied postmergence at 250 to 375 pounds per acre, will control the following weeds in non-cropland areas (Table 5). Use the higher labeled rates to extend length of control. Use the higher rates on sites with fine soil textures and on sites with more than 2% organic matter.

**Weeds Controlled from Preemergence Application to Bare Ground Prior to Weed Seed germination**  
 Table 5

Common Name	Scientific Name	Common Name	Scientific Name
Barnyardgrass	(Echinochloa crus-galli)	Lambsquarters,	(Chenopodium album)
Bluegrass annual	(Poa annua)	common	
Carpetweed	(Mollugo verticillata)	Lovegrass	(Eragrostis spp.)
Carolina Geranium	(Geranium carolinianum)	Nutsedge, yellow	(Cyperus esculentus)
Chickweed, common	(Stellaria media)	Nutsedge, purple	(Cyperus rotundus)
Chickweed,mousear	(Cerastium vulgatum)	Panicum(Texas,Fall)	(Panicum spp.)
	(from seed)	Pigweed, Redroot	(Amaranthus retroflexus)
Crabgrass	(Digitaria spp.)	Purslane, Common	(Portulaca oleracea)
(large, Smooth) <sup>3</sup>		Pusley, Florida	(Richardia scabra L.)
Crowfootgrass	(Dactyloctenium aegyptium.)	Sedge, globe	(Cyperus globulosus)
Cupgrass, Woolly	(Eriochloa villosa)	Sedge, cylindrical	(Cyperus retrorsus)
Foxtails, Annual	(Setaria spp..)	Sedge, Surinam	(Cyperus surinamensis)
Goosegrass	(Eleusine indica)	Sedge, Texas	(Cyperus polystachyos)
Hairy Bittercress	(Cardamine hirsute)	Shephardspurse	(Capsella bursa-pastoris)
Henbit	(Lamium amplexicaule )	Signalgrass, broadleaf	(Bracharia playphylla)
Itchgrass	(Rottboellia exaltata)	Speedwell	(Veronica spp.)
Knotted, prostrate	(Polygonum aviculare)	Sprangletop	(Leptochola spp.)
Kochia	(Kochia scoparia)	Spurge, prostrate	(Euphorbia humistrata)
Kyllinga, green	(Kyllinga brevifolia)	Witchgrass	(Panicum capillare L.)
Kyllinga, false green	(Kyllinga gracillima)	Woodsorrel, yellow	(Oxalis stricta).
		(from Seed)	

SPREADER SETTINGS						
lbs/ 1000 sq.ft.	Cone setting	2.0 lbs	3.0 lbs	4.0 lbs	5.0 lbs	6.0 lbs
Accu-Pro 2000	6	K 1/2	L 1/2	M 1/2	N 1/2	O 1/2
SR 2000	6	K 1/2	L 1/2	N	O 1/2	P
Lesco 21820	0.25 Closed	C 1/2	D 1/2	E 1/2	F 1/2	G 1/2
SpykerR 76-22		4.1	4.6	4.5	4.7	4.9
PSB Bigfoot	A.5	J	J 1/2	K 1/2	L 1/2	M
Lely HR	II-B	4.75	5.25	6	6.25	7
Vison 203		13	15	17	19	21

**CONDITIONS OF SALES AND LIMITATIONS OF WARRANTY AND LIABILITY**  
**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.