



Howard Johnson's

# OXADIAZON 1.0 PLUS

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF MANAGEMENT & LANDSCAPE PERSONNEL NOT FOR USE ON HOME LAWNS. FOR USE ON GOLF COURSES, NURSERIES, PARKS, SOD FARMS, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS.

A Turf Fertilizer plus pre-emergent weed control product. For use on varieties of Perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustine grass, Seashore Paspalum, Tall Fescue and Zoysiagrass. Controls Crabgrass, Goosegrass, Poa annua, Annual Sedge and other weeds as listed on Golf Courses, Parks and recreational and Ornamental turf.

### ACTIVE INGREDIENT:

Oxadiazon\*: ..... 1.0%  
OTHER INGREDIENTS: ..... 99.0%  
TOTAL: ..... 100.0%

\*CAS [19666-30-9]

EPA Reg. No. 53883-180-32802

EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (C)

Manufactured for: Howard Johnson's Enterprises, Inc.  
9675 S 60th Street, Franklin, WI 53132

## GUARANTEED ANALYSIS

### 15-0-10

Total Nitrogen (N) 15%  
15.00% Urea Nitrogen\*  
Soluble Potash (K<sub>2</sub>O) 10%

Derived From: Muriate of Potash, Polymer Coated Urea, Urea

\* 6.00% Slowly Available Nitrogen from Polymer Coated Urea

Chlorine, (Cl), not more than 6.00%

F315

Bulk Density 55 Item Number FGP54367  
100206647

### Net Weight 50 lbs (22.70 kg)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

## KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust, or contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):** Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; chemical resistant gloves; chemical resistant footwear plus socks; protective eyewear; headgear for overhead exposure; chemical-resistant apron when cleaning equipment. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS:** Applicators should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### FIRST AID

**IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.  
**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.  
**IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.  
**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Apply only as specified on this label.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls; chemical resistant (Barrier Laminate) gloves, chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NQI within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, and nurseries. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse. Do not allow entry to areas treated until dusts have settled or if watering in is required, until treated area and grass is dry.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE AND SPILL PROCEDURES:** Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

**CONTAINER DISPOSAL:** If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Do not rinse unless required for recycling. If recycling is not available: then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. If partly filled: Call your local solid waste agency for disposal instructions. Never pour unused product down any indoor or outdoor drain.

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

### WEEDS CONTROLLED:

Annual Bluegrass	Annual Sedge	*Barnyardgrass	Bittercress	Carpetgrass	Common Groundsel
Common Purslane	Crabgrass	*Evening Primrose	Fireweed	Field Sandbur	
Foxtail, Green, Bristly	Fall Panicum	*Florida Pusley	*Galinsoga	Golden Ragwort	*Goosegrass
**Jungle Ricegrass	Lambsquarter	**Liverwort	Oxalis	**Panama Paspalum	Pennsylvania Smartweed
Redroot Pigweed	Sensitive Plant	Shepherdspurse	Southistle	Speedwell-Veronica	Spotted Catsear
Spurge; Garden, Petty, **Prostrate	*Stinging Nettle	*Swinecress	Virginia Buttonweed	Yellow Woodsorrel	

\*For these weeds use 3 to 4 lbs. A.I./acre (equal to 300 to 400 lbs of this product per acre or 6.9 lbs. to 9.2 lbs. of this product per 1,000 sq.ft.).

\*\*For these difficult-to-control weeds use 4 lbs. A.I./acre (equal to 400 lbs. of this product/acre or 9.2 lbs. per 1000 sq.ft.).

**WHERE, HOW MUCH AND WHEN TO USE ON TURFGRASS:** This product may be used at full rate or as a "split" application in established Perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf, at apartment/condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, cemeteries and other ornamental and recreational turfgrass areas. Not for use on home lawns. The turf should be free of leaves and thatch and in condition to allow the granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply with a granule spreader. Do not apply to turf when it is wet. When using at full rate do not overlap spreader applications, especially on St. Augustinegrass since discoloration "streaking" may occur from the "doubled-up rate." See section on this label entitled PRECAUTIONS IN USING ON TURFGRASS for maximum rate for St. Augustinegrass and Bentgrass of 3 lbs. A.I. per acre which is equivalent to 6.9 lbs of this product per 1000 sq. ft. After applications if rain is not expected within 24 hours sprinkler irrigate the treated turf with up to a 1/4 inch of water to release the herbicide for maximum weed control and quicker response from the fertilizer nutrients. This product is a pre-emergence herbicide and must be applied prior to weed seed germination. It will not control growing weeds.

\*See paragraph below for instructions to apply the 3 lb. or 4 lb. rate as a split application.

**PRECAUTIONS IN USING ON TURFGRASS:** For St. Augustinegrass and Bentgrass turf, **DO NOT** exceed the rate of 3 lbs. active ingredient per acre which is equal to 6.9 lbs. of this product per 1000 sq. ft. **DO NOT** apply to **Bentgrass** turf mowed less than 3/8 inch. This product may cause discoloration of **Bentgrass**, **Bermudagrass** and **St. Augustinegrass** but this discoloration is temporary. Avoid contact with flowers and shrubs except as recommended elsewhere for ornamental plants on this label. **DO NOT** apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. **DO NOT** apply to **Dichondra** or **Centipede**grass. **DO NOT** use on golf course putting greens or tees. Note: For newly seeded lawns **DO NOT** apply this product until after the second or third cutting. For best turf management, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

**SPRING AND SPLIT APPLICATION ON GOLF COURSES, The 3 LB. & 4 LB. per acre rate can be applied as a split application.**


Weed control and better turf management can be programmed by splitting the per acre rate of active ingredient into "two" separate applications. For example, to accomplish this in region #5 (see map of US) apply 1.5 lbs. A.I. (for a required 3 lbs. A.I./acre rate) or 2 lbs. A.I. (for a required 4 lbs. A.I./acre rate) as the first application 1 to 2 weeks before estimated weed seed germination or in late December or early January. Follow with the second "1/2 rate" in 60 days or the first of March. The timing for these applications should be closely programmed to the geographic region (see map of US) and adjusted to an "early" or "late" spring. Do not split the 2 lb. rate. The 3 lb. rate **should not be split** if applied to new turf areas first coming under turf management where weed pressure is expected to be heavy.

### WHERE, HOW MUCH, AND WHEN TO USE, AROUND ORNAMENTALS

Can be applied anytime of the year prior to weed seed germination to both newly transplanted and established actively growing or dormant ornamental plants where both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control activity. Do not mix into soil. Disturbing or cultivating the soil after treatment may reduce weed control. Do not apply granules to wet foliage or under conditions in which granules will collect on leaves, if so, sprinkler-irrigate the plants to move granules off foliage and down to soil level. Do not apply in greenhouses as plant injury may likely result. Use the higher application rates for areas that have a history of heavy weed pressure. For best weed control around ornamentals first remove any existing weed growth and if possible rake, cultivate and smooth the soil surface by breaking-up clumps. Broadcast the granules evenly over the prepared soil by applying the 2 to 4 lbs a.i. per acre, equal to 200 to 400 lbs. of this product / acre or 4.6 to 9.2 lbs. per 1000 sq.ft.

### CRABGRASS SEED GERMINATION DATES

Hawaii included in January 1 to March 20 Zone



### SPREADER SETTINGS FOR POUNDS OF PRODUCT PER ACRE

The following spreader settings are for new equipment. Used, or heavily worn equipment may require adjustments. For proper setting calibration refer to the spreader manufacturer's Maintenance Manual. Do not apply this product until equipment is properly calibrated to accurately apply the desired lbs. of product/acre, as recommended on this label.

SPREADER SETTINGS	1.5 lb. A.I./acre				2 lb. A.I./acre				3 lb. A.I./acre				4 lb. A.I./acre			
	4.0				4.3				5.0				4.3 x2			
Spyker 24-22 Rotary	12 (F)				13 (F-G)				16 (H)				(F-G) x2			
Lesco Rotary	4.4				4.6				6.0				4.6 x2			
Prizelawn CB Rotary	17				17				18				20			
EarthWay 2400 Rotary	18 x2				18 x2				18 x2				18 x2			
Cyclone Pro	3.75				4.25				5.0				4.25 x2			
Scotts R8A	J				J				L				J x2			
Lely Broadcast	2.8				3.2				3.8				3.2 x2			
Bigfoot	H				I				J				I x2			
Cub Cadat	14				15				18				N/A			

1.0 Oxadiazon 53883-180 CSI08-26-09(hje 12-05-15)

### PRE-EMERGENCE WEED CONTROL (Refer to map of US)

Weeds Controlled	lbs A.I. per acre	lbs/product 1000 sq ft
Crabgrass ( <i>Digitaria</i> spp.)	2 to 4 lbs. A.I. / acre	4.6 to 9.2 lbs. / 1000 sq. ft. (200 to 400 lbs. / acre)
Goosegrass ( <i>Elymus indica</i> )		
Field Sandbur ( <i>Cenchrus insuetus</i> )		
Annual Sedge ( <i>Opuntia compressa</i> )		
Directions: Apply in late winter or early spring prior to weed seed germination. See map. In Areas 4 & 5 application made after April or May may not be fully effective.		
Annual Bluegrass ( <i>Poa annua</i> )	2 to 4 lbs. A.I. / acre	4.6 to 9.2 lbs. / 1000 sq. ft. (200 to 400 lbs. / acre)
(See note below)		
Directions: Apply in late summer to early fall prior to weed seed germination.		
Virginia Buttonweed ( <i>Diodia virginiana</i> )	4 lbs. A.I. / acre	9.2 lbs. / 1000 sq. ft. (400 lbs. / acre)
Pigweed ( <i>Amaranthus</i> spp.)		
Florida Pusley ( <i>Richardia scabra</i> )		
Oxalis ( <i>Oxalis</i> spp.)		
Stinging Nettle ( <i>Urtica dioica</i> )		
Carpetgrass ( <i>Axonopus officinis</i> )		
Directions: Apply in early spring prior to weed seed germination.		

### FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF

Crabgrass, Goosegrass and other annual weeds as listed	3 to 4 lbs. A.I. / acre	6.9 to 9.2 lbs. / 1000 sq. ft. (300 to 400 lbs. / acre)
Directions: For use in Bluegrass, Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup.		

### ANNUAL BLUEGRASS CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED WITH PERENNIAL RYEGRASS ONLY

Annual Bluegrass ( <i>Poa annua</i> spp.)	2 lbs. A.I. / acre	4.6 lbs. / 1000 sq. ft. (200 lbs. / acre)
Directions: Apply in late summer to early fall prior to <i>Poa annua</i> spp. seed germination. Any <i>Poa annua</i> spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where <i>Poa trivialis</i> is present in the overseed mixture. Apply the full 2 lbs. A.I. rate in one application. Note: In many areas Annual Bluegrass exists as at least two biotypes or subspecies of <i>Poa annua</i> . The true annual biotype, " <i>Poa annua</i> spp. <i>annua</i> " which germinates from seed every year is controlled while the biotype " <i>Poa annua</i> spp. <i>raptans</i> " which behaves like a perennial and survives through the summer as an established plant cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning your locale.		

### ANNUAL WEED CONTROL DURING ESTABLISHMENT FROM "SPRINGS" OF BERMUDAGRASS, ZOYSIAGRASS, AND SEASHORE PASPALUM

Crabgrass, Goosegrass, Annual Sedge and other annual weeds	2 to 3 lbs. A.I. / acre	4.6 to 6.9 lbs. / 1000 sq. ft. (200 to 300 lbs. / acre)
Refer to Map of U.S. on this label. For Bermudagrass make application immediately prior to, or, immediately after sprigging. For Seashore Paspalum and Zoysia make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" areas with up to 1/4 inch water after application.		

### READ ENTIRE LABEL BEFORE USING

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use of application, all of which are beyond the control of Howard Johnson's. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Howard Johnson's is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Howard Johnson's disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product. **LIMITATION OF WARRANTIES:** The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Howard Johnson's option, the replacement of the product.